Supplementary Table 3. Musculoskeletal representation of participants (n=378)

Variables	n (%) median (Q1 – Q3)	Scale scores	
		Median (Q1-Q3) r_s	p*
P-NRS score	5 (4 – 6)	0.609	**<0.001**
Regions of pain*			
Shoulder	119 (31.5)	55 (42 – 66)	**0.045**
Elbow	25 (6.6)	55 (42 – 65)	0.598
Wrist	57 (15.1)	55 (41 – 66)	0.243
Hand	30 (7.9)	51 (45 – 65)	0.268
Arm	44 (11.6)	57 (46 – 68)	**0.014**
Forearm	19 (5.0)	54 (42 – 66)	0.368
Hip	63 (16.7)	65 (50 – 76)	**<0.001**
Knee	87 (23.0)	59 (41 – 71)	**0.009**
Ankle	45 (11.9)	65 (54 – 73)	**<0.001**
Foot	46 (12.2)	61 (49 – 73)	**0.003**
Thigh	26 (6.9)	63 (51 – 83)	**0.006**
Leg	65 (17.2)	64 (40 – 74)	**<0.001**
Head	52 (13.8)	57 (42 – 68)	0.090
Neck	182 (48.1)	54 (42 – 66)	**0.003**
Back	168 (44.4)	54 (42 – 67)	**0.015**
Low back	193 (51.1)	55 (42 – 68)	**<0.001**
Place where pain first felt:			
At workplace	81 (21.4)	53 (43 – 64)	
At home	140 (37.0)	52 (36 – 65)	
Outside and outdoor	26 (7.0)	51 (32 – 71)	0.562
Outside and indoor	10 (2.6)	41 (33 – 63)	
Don’t remember	121 (32.0)	52 (41 – 64)	
Pain’s relation to work	210 (55.6)	52 (39 – 64)	0.897
Consulting to a physician	227 (60.1)	58 (42 – 69)	**<0.001**
Medical certificate (n=378)	84 (22.2)	66 (53 –76)	**<0.001**
Medical certificate Days (n=62)	10 (5 – 30)	0.081	0.534

Data were summarized as median (Quartile 1- Quartile 3), frequency (percentage), or Spearman correlation coefficient.

*The Wilcoxon rank-sum test or Spearman rho correlation coefficient test results were given based on the independent variable type.